



Relationships Between Profiles of Physical Activity and Major Mobility Disability in the LIFE Study

Jason Fanning, PhD,* W. Jack Rejeski, PhD,* Shyh-Huei Chen, PhD,[†] Jack Guralnik, MD,[‡] Marco Pahor, MD,[§] and Michael E. Miller, PhD[†]

OBJECTIVES: To examine the relationship between time spent in light physical activity (LPA) and in moderate to vigorous physical activity (MVPA) and the pattern of accumulation on the risk for major mobility disability (MMD) in a large multicenter study of physical activity (PA) and aging, the Lifestyle Interventions and Independence for Elders (LIFE) study.

DESIGN: Data were collected from individuals randomized to a PA intervention as part of the LIFE study, an eight-center single-blind randomized clinical trial conducted between February 2010 and December 2013.

SETTING: Lifestyle Interventions and Independence for Elders Study

PARTICIPANTS: Older adult participants (78.4 years; N = 507) at risk for MMD.

INTERVENTION: All older adults included in these analyses were randomized to a structured PA intervention that included two center-based plus three to four home-based exercise sessions per week with a primary goal of walking for 150 minutes weekly. Participants attended the intervention for 2.5 years on average.

MEASUREMENTS: MMD was defined as the inability to complete a 400-m walk within 15 minutes and without assistance. Physical function was assessed via the Short Physical Performance Battery (SPPB). Actigraph accelerometers were used to quantify amount and variability in LPA and MVPA.

RESULTS: In an adjusted Cox proportional hazards regression, we identified a significant interaction ($P = .017$)

between SPPB score and LPA amount and variability such that more LPA was associated with a reduced risk for MMD among those with higher initial function, as was lower variability (eg, via distributing LPA across the day). The SPPB \times MVPA interaction was significant ($P = .04$), such that more MVPA was associated with lower MMD risk among those with lower function. Finally, greater MVPA variability was associated with lower risk for MMD.

CONCLUSION: A prescription of PA for older adults should account for key factors such as physical function and emphasize both amount and pattern of accumulation of PA from across the intensity continuum. *J Am Geriatr Soc* 00:1-8, 2020.

Keywords: physical activity; sedentary behavior; mobility; function

From the *Department of Health and Exercise Sciences, Wake Forest University, Winston-Salem, North Carolina; [†]Department of Biostatistical and Data Sciences, Wake Forest School of Medicine, Winston-Salem, North Carolina; [‡]Department of Epidemiology and Public Health, University of Maryland School of Medicine, Baltimore, Maryland; and the [§]Department of Aging and Geriatric Research, University of Florida, Gainesville, Florida.

Address correspondence to Jason Fanning, PhD, Worrell Professional Center 2164B, PO Box 7868, Winston-Salem, NC 27109.
E-mail: fanninj@wfu.edu

Trial Registration clinicaltrials.gov Identifier: NCT01072500

DOI: 10.1111/jgs.16386

Impaired mobility, a key marker of physical disability, is critical to the loss of functional independence as people age.^{1,2} More than 2 decades ago, reviews concluded that physical inactivity was the strongest single predictor of physical disability in aging.^{3,4} Conventionally, physical activity (PA) research has focused on structured exercise, or discrete bouts of purposeful moderate to vigorous intensity physical activity (MVPA) (>3.0 metabolic equivalents [METs]). Although the health benefits of regular MVPA are well established,⁵ interest in light physical activity (LPA) has increased in recent years. This is particularly important for older adults because LPA may be less aversive and more attainable than more intense activity, especially for those who are frail or pre-frail.^{6,7} The purpose of the present study was to examine the relationship between time spent in LPA and MVPA and the pattern of accumulating these activities on the risk for major mobility disability (MMD) among older adults with compromised physical function

who participated in the Lifestyle Interventions and Independence for Elders (LIFE) study.⁸

PA is beneficial in improving mobility-related physical disability,^{9,10} with a number of randomized controlled trials (RCTs) showing that structured exercise improves mobility, even among older adults with severe chronic health conditions.¹¹⁻¹³ As such, the health-promoting effects of MVPA are endorsed by institutions such as the American College of Sports Medicine¹⁴ and the World Health Organization.¹⁵ In 2014, Pahor and colleagues published the results of the LIFE study, a multicenter RCT comparing a PA program involving the promotion of MVPA with a health education (HE) intervention on MMD among older adults who were frail or pre-frail. They found that participants in PA had a lower 2-year incidence of MMD than those in HE.⁸ A subsequent publication reported that those participants classified as achieving the highest quartile for change in MVPA had the greatest reduction in risk for MMD. Interestingly, the lower bound of MVPA duration for the fourth quartile was quite modest at just 43 minutes per week.¹⁶

In recent years, interest has been increasing in PA that falls below the threshold for MVPA. For example, whereas the first edition of the PA Guidelines for Americans recommended between 150 and 300 minutes per week of MVPA lasting at least 10 minutes per bout, the recent second edition encourages increasing both MVPA and LPA. Of note is that within the LIFE study, we observed that as many as 25% of participants were unable to reach the accelerometry-defined threshold for MVPA during center-based exercise sessions,⁷ and they often had to stop and rest or were unable to sustain the MVPA threshold for an extended period of time; their activity was highly variable. Moreover, the second edition of the guidelines removed the recommendation that activity persist for at least 10 minutes, reflecting a growing interest in the effect of not only the amount of PA on health but also the pattern in which it is accumulated. For instance, does participating in LPA or MVPA frequently throughout the day have different health effects as compared with fewer, longer bouts of the behaviors?

The objective of the current study was to examine the relationships between the amount and variability in both MVPA and LPA on subsequent risk for MMD among at-risk older adults who participated in the PA intervention of the LIFE study.⁸ We first examined relationships between the amount and variability in MVPA with MMD. We hypothesized a dose-response effect indicating that participating in more MVPA would be associated with lower risk for MMD, although this effect would be small because all individuals participated in a program that targeted increasing MVPA. We also expected that greater variability in minutes per hour of MVPA (reflecting fewer, longer bouts of MVPA relative to lower variability) would be associated with reduced risk. Regarding LPA, we hypothesized that greater volume would be associated with lower risk for MMD and that, across individuals, a more distributed pattern of movement (ie, lower variability) would be associated with lower risk. Lastly, we explored potential interactions between differences in basic levels of physical functioning as assessed by the Short Physical Performance Battery (SPPB) with metrics for MVPA and LPA, because previous publications suggested that lower functioning individuals

benefit most from the intervention,⁸ engage in high levels of sustained sedentary behavior,¹⁷ and are uniquely challenged in their ability to meet accelerometer cut points for MVPA specific for older adults.⁷

METHODS

Design and Participants

Data were collected from individuals randomized to a PA intervention as part of the LIFE study, an eight-center single-blind RCT that was conducted between February 2010 and December 2013 (detailed procedures are described in Fielding et al¹⁸). LIFE examined the effect of a long-term structured exercise intervention on transition to MMD (ie, inability to walk 400 m) among those able to walk 400 m at baseline. Participants were required to have an SPPB score of 9 or lower of 12 (45% had SPPB <8). Eligible individuals were 70 to 89 years of age, low to active (<20 min/wk of structured PA; <125 min/wk MVPA over last month), able to walk 400 m in less than 15 minutes, without major cognitive impairment, and able to participate safely in the PA intervention as determined by medical history, electrocardiogram, and physical examination. The PA intervention was well tolerated, as supported by a median attendance rate of 71% across nearly 2 years including medical leave. Additionally, those in the intervention condition engaged in an additional 40 minutes per week of MVPA at month 6 that was maintained at 12 and 24 months.⁸

Randomization

Participants were randomized to receive PA or a HE control group, and randomization was stratified by field center and participant sex. All participants received an orientation appointment with a health educator, at which point they reviewed their randomization assignment and clarified any outstanding questions. Because our goal in the current study was to examine how amount and variability of MVPA and LPA relate to risk for MMD, we restricted our analyses to participants randomized to the PA intervention.

PA Intervention

The focus of the PA intervention was on aerobic PA via walking, with a goal of 150 weekly minutes of MVPA paired with brief balance, flexibility, and lower body strengthening exercises. Participants aimed to engage in exercise 5 to 6 days per week including two weekly center-based exercise sessions. For the first 2 to 3 weeks of training, participants progressed from exercising at a light intensity to a rating of perceived exertion¹⁹ of 13 out of 20 during walking, and 15 to 16 out of 20 during strength training (ClinicalTrials.gov identifier NCT01072500).¹⁸

Measures

Major Mobility Disability

Our dichotomous outcome measure was onset of MMD, defined as the inability to complete a 400-m walk test

within 15 minutes without assistance. This test was completed before randomization and again at 6-month intervals.

Actigraphy

Participants wore a GT3X accelerometer (Actigraph, Pensacola, FL) on the right hip during waking hours (except while bathing or swimming) for 1 week at baseline and at the 6-, 12-, and 24-month visits. Data were processed in ActiLife (v.4.4.1), using the Choi²⁰ algorithm to classify non-wear time. Specifically, periods of 90 consecutive minutes with zero counts per minute (CPM), allowing for a 1-minute spike tolerance, were classified as non-wear. Participants with at least 3 days of at least 600 minutes of wear were retained in analyses. Sedentary was defined as less than 100 CPM, LPA was defined as 100 to 1041 CPM, and MVPA was defined as more than 1040 CPM.²¹

Because low-functioning older adults often achieve MVPA at intensities below conventional accelerometer cut points for MVPA, we ran an additional sensitivity analysis in which 100 to 760 represented LPA and more than 760 classified MVPA.^{7,22,23} Average daily minutes of LPA and MVPA were computed across valid days. Daily variance in minutes per hours of LPA and MVPA were computed and averaged across valid days before a square root transformation was applied.²⁴ This measure provides insight into the distribution of movement during the day, as illustrated in Figure 1 where the profile marked with the solid line and the profile denoted by the dashed line represent different LIFE participants with similar minutes of daily LPA (solid = 7.97 min/h; dashed = 8.01 min/h), but relatively high (solid = 9.87) and low (dashed = 3.93) variability in LPA. Here, profile A (high variability) engages in fewer but longer bouts of LPA, with lengthy daily periods with little LPA, and profile B (low variability) frequently maintains approximately 5 to 20 minutes per hour of LPA during waking hours.




Figure 1. Two individuals with similar average daily minutes of light physical activity (LPA) and high (solid line) vs low (dashed line) LPA variability.

Analyses

We explored relationships between amount and variability in both MVPA and LPA and risk of MMD using a series of Cox proportional hazards regression models that incorporated these activity measures as time-dependent covariates (ie, updating the accelerometry variables at 6-, 12- and 24-mo intervals) using time until the initial occurrence of MMD as the dependent variable. Each model controlled for baseline sex (as a stratifying factor for the baseline hazard), age, body mass index (BMI), race, history of high blood pressure, cardiovascular disease, lung disease, and osteoarthritis. We first fit a model for MVPA that included the three-way and lower order interaction terms between baseline SPPB classified as frail (scores ≤ 7) or pre-frail (scores 8-9),²⁴ amount and variability of MVPA, and all covariates. The highest order interaction terms with $P < .05$ via likelihood ratio tests, and associated lower order interactions, were retained. Next, this model-building process was repeated for LPA, log-transformed due to skewness. In our final model we included all previously identified LPA and MVPA amount and variability terms with $P < .05$.

RESULTS

Of the 14 831 individuals assessed for eligibility, 818 were randomized to PA (Figure 2; participant description provided in Table 1). Following accelerometer data cleaning procedures, 507 individuals had accelerometer data at baseline (7.5 d on average) and follow-up MMD data and were included in the analyses. Of these, updated activity information was obtained at month 6 on 354 participants (6.6 d on average), 406 at month 12 (6.9 d on average), and 382 at month 24 (6.9 d on average). Baseline participant characteristics of this group were previously reported (Fanning et al,²⁴ Table 1 of citation²⁴) and reveal that the average age and BMI were 78.4 years and 30.4 kg/m², respectively, with 64% women and 26% from minority racial groups. Included participants had a median follow-up time for MMD (time until censoring or an event) of 2.5 years, 291 were categorized as pre-frail, and 144 were determined to have MMD at least once during follow-up.

Results for all Cox proportional hazards models are provided in Table S1. In the preliminary model (amount and variability of MVPA, SPPB frailty classification), there was no significant three-way interaction ($P = .64$). After removal of nonsignificant ($P > .05$) higher order interactions, there remained a two-way interaction between the amount of MVPA and baseline SPPB ($P = .017$), and a main effect for variability ($P = .058$; hazard ratio [HR] = .81; 95% confidence interval [CI] = .64-1.00) of borderline statistical significance. For frail individuals, more MVPA was associated with a lower risk of MMD (HR = .61 for a 1-min/h difference; 95% CI = .31-1.08); whereas among pre-frail individuals, more MVPA was associated with an elevated risk (HR = 1.12 for a 1-min/h difference; 95% CI = .70-1.70). Our model that included main effects for amount and variability in LPA, SPPB classification, and all relevant interaction terms revealed a significant three-way interaction term ($P = .003$) that we explored further in the final full model




Figure 2. Participant flow. ¹Participants who did not receive the allocated intervention (ie, attended no intervention sessions). ²Discontinuation of the intervention was operationalized as participants who did not attend at least one intervention session during their last 6 months of follow-up before the last planned follow-up visit date. Deaths and intervention withdrawals are included in these numbers. MMD, major mobility disability.

Table 1. Participant Characteristics

| Baseline characteristic | Event classification | | Comparison of event groups P value | Overall (n = 507) |
|---|------------------------|---------------------|---------------------------------------|-------------------|
| | No MMD event (n = 363) | MMD event (n = 144) | | |
| Age, y, mean (SD) | 78.1 (5.09) | 79.4 (5.74) | .01 | 78.4 (5.31) |
| Sex, female, n (%) | 234 (64.5) | 92 (63.9) | .90 | 326 (64.3) |
| Race/ethnicity, non-Hispanic white, n (%) | 26 (72.7) | 109 (75.7) | .26 | 373 (73.6) |
| Education, some college, n (%) | 248 (68.3) | 82 (56.9) | .04 | 330 (65.1) |
| SPPB total score, mean (SD) | 7.7 (1.45) | 6.9 (1.66) | <.001 | 7.47 (1.56) |
| SPPB ≤7, yes, n (%) | 133 (36.6) | 83 (57.6) | <.001 | 215 (42.6) |
| 400 m walk speed, m/s, mean (SD) | .87 (.15) | .74 (.16) | <.001 | .83 (.16) |
| BMI, mean (SD) | 30.1 (5.52) | 31.0 (6.70) | .12 | 30.4 (5.89) |
| History of hypertension, yes, n (%) | 247 (68.6) | 103 (72.5) | .39 | 350 (69.7) |
| History of CVD, yes, n (%) | 87 (24.0) | 58 (40.3) | <.001 | 145 (28.6) |
| History of lung disease; yes, n (%) | 65 (18.1) | 16 (11.1) | .06 | 81 (16.1) |
| History of arthritis, yes, n (%) | 62 (17.2) | 25 (17.4) | .96 | 87 (17.2) |

Abbreviations: BMI, body mass index; CVD, cardiovascular disease; MMD, major mobility disability; SPPB, Short Physical Performance Battery.

The final model included amount and variability of MVPA plus their interaction, as well as LPA amount, variability, amount \times variability, and SPPB \times amount \times variability. Here, conclusions from the MVPA amount \times SPPB interaction and variability terms were essentially unchanged, and results for LPA were similar to the model without MVPA. Specifically, the baseline SPPB \times LPA amount \times LPA variability interaction retained statistical significance ($P = .02$). This interaction is depicted in Figure 3, with the upper panel depicting frail individuals and the lower panel depicting pre-frail individuals. LPA amount is depicted on the x-axis and variability on the y-axis.

All plots provide HRs (obtained by back-transforming the log-transformed LPA values) for a 1-unit difference on each axis at selected values of the other axis (eg, a 1 min/h




Figure 3. Difference in risk for major mobility disability of a 1-unit higher level of variability at different amounts of light physical activity (LPA) (vertical lines). The figure also shows the difference in risk associated with a 1 min/h higher level of amount at different levels of variability in LPA (horizontal lines) among those with a Short Physical Performance Battery (SPPB) score of 7 or lower (bottom panel) and an SPPB score of 8 to 9, extracted from a model including all activity predictors. Variability hazard ratios (HRs) are shown at selected values of 6 vs 5, 9 vs 8, and 11 vs 10; number of HRs are shown at selected values of 7 vs 6 min/h, 13 vs 12 min/h, and 17 vs 16 min/h. Red dots represent activity measurement intervals followed by MMD events. HRs for LPA were obtained through back-transformation from models generated using log-transformed values of LPA.

difference in amount at a specific level of variability). As depicted in the interaction plot, for frail participants, both amount and variability in LPA were minimally related to risk for MMD. Among pre-frail participants who were engaging in very little LPA, there was a risk reduction for greater amount and variability of LPA. At higher amounts of LPA, adding more movement had little effect on risk. However, for these higher active, higher functioning participants, lower variability was associated with a risk reduction. Finally, our sensitivity analyses using a lower threshold for MVPA (ie, 760 CPM) did not differ meaningfully from the results presented here, except that findings related to variability in higher functioning individuals appeared amplified (Figure S1 and Table S2).

DISCUSSION

It is important to place our results in the context of the LIFE sample and PA intervention. Participants were 70 to 89 years of age, obese, and physically compromised (SPPB ≤ 9 ; average gait speed during 400-m walk of .83 m/s) at baseline. They had multiple comorbidities, and none had major cognitive impairment. The structured PA sessions were individualized and progressed toward a goal of 30 minutes of daily walking at a moderate intensity, 10 minutes of primarily lower extremity strength training by means of ankle weights, and 10 minutes of balance training/large muscle group flexibility exercises.¹⁸ Notably, only 2.5% of participants in the LIFE study were able to reach a common cut point specific for older adults for MVPA during the adoption phase of training.⁷ Moreover, many participants had to take a break during exercise.⁶ Thus when older adults who are either frail or pre-frail exercise, they frequently engage in what clinicians would traditionally conceptualize as LPA or non-exercise PA (ie, brief bouts of oftentimes slow walking).²⁵

We would like to emphasize here that even this movement that appears low in intensity is likely to fall within the individual's true moderate to vigorous range (ie, >3.0 METs). Older adults with limited function may achieve 3.0 METs at walking speeds as slow as 1.5 mph that falls well below what is characterized as LPA in the current PA guidelines.^{22,25,26} We illustrate this point in Figure 4, which shows changes in CPM the first 6 months of the intervention by SPPB category. Those with lower SPPB scores increased activity in lower intensity bouts compared with those with higher function.

Finally, older adults are prone to serious health events. Within the PA group of the LIFE study, 48.4% of participants had a hospitalization unrelated to the intervention. A total of 58.6% went on medical leave at least once with a median length of 49 days (interquartile range [IQR] = 21-140 d).⁸ Mobility was often limited following this leave, and interventionists had to focus on strength and balance to assist the participant in returning to extended bouts of walking. Only 14.5% (5.9% due to death) of participants dropped out or were unable to continue with the intervention and complete the final assessments. In sum, these were resilient older adults challenged by compromised function and comorbidities, engaging in an intervention composed of aerobic MVPA paired with LPA and strength and balance training as needed.




Figure 4. Change in accelerometer counts per minute (CPM) (median difference per day) in the first 6 months of the intervention by the Short Physical Performance Battery (SPPB) score.

Here we investigated the extent to which the amount and variability in both MVPA and LPA were associated with the development of MMD. Our results made clear that the relationship between PA and MMD must be viewed through the lens of function. What follows is a discussion of findings for pre-frail (ie, $SPPB = 8-9$) and frail (ie, $SPPB \leq 7$) individuals.

Structuring Movement Interventions for Pre-Frail Older Adults

Data from the National Health and Aging Trends study ($N = 4377$)²⁷ indicate that as many as 14.6% of older adults aged 65 to 69 years are pre-frail ($SPPB = 8-9$). Functional scores decrease with age such that 77.2% of those aged 90 or older qualify as frail, and an additional 12% are pre-frail. For pre-frail individuals, “move more, more often” appears a good recommendation for reducing risk for MMD. Interestingly, counter to our hypothesis and to an analysis of the overall study sample,¹⁶ additional minutes of MVPA were associated with a small increase in risk for MMD, a phenomenon described in prior LIFE study publications and perhaps attributable to increased risk for injury.⁸ Both pre-frail and frail participants demonstrated benefit from more sustained bouts (ie, higher variability) of MVPA, suggesting those most capable of adhering to the intervention had a generally lower risk for MMD.

In line with our hypothesis, greater LPA was associated with lower risk for MMD among members of this group with very low levels of LPA, an effect that held when controlling for participation in MVPA. Thus for pre-frail individuals who engage in low levels of LPA, clinical recommendations should first emphasize “moving more” to increase LPA. Recommended activities might include a variety of lifestyle behaviors such as frequent leisure walks and engaging in hobbies that challenge a variety of functional abilities. Once a greater total volume of LPA has been achieved, recommendations can be made to “move more often” by engaging in frequent bouts of LPA and MVPA throughout the day to maintain a high volume of movement

with reduced variability. In sum, pre-frail older adults are likely to benefit from recommendations emphasizing greater volume in movement throughout the day paired with daily participation in MVPA.

Structuring Movement Interventions for Frail Older Adults

Our findings for frail participants highlighted unique challenges for both the assessment and promotion of health-promoting PA among this group. Nationally, less than one-third of older adults younger than 80 years fall in this category, but this number rises quickly with increasing age.²⁷ Poorer functional status places individuals at a greater risk for mobility disability and acts as a powerful barrier to engaging in MVPA.²⁸ Thus a key first step in exercise prescription for frail older adults should be the development of strength and balance to support and foster improved mobility.²⁹ Our experience in the LIFE study following return from medical leave due to acute health events supports this position. Then as older individuals find they are more capable of MVPA, recommendations should focus on accumulating a larger daily volume of MVPA, recognizing that we are talking about a low criterion for defining MVPA. It is intriguing to note there were no MMD events observable in individuals engaging in very high levels of LPA (Figure 3). It may be that older adults with a high degree of functional impairment benefit from a high volume of this behavior.

Integrating Function-Oriented Prescriptions into a Movement Medicine Framework

Our results add to a building “movement medicine” approach to PA prescription for older adults, one that considers the individual’s specific needs and desired outcomes when crafting clinical activity recommendations. Unfortunately, there are numerous barriers to prescribing PA in the clinic including lack of time and inadequate training on the complexities of traditional approaches to activity prescription

that has resulted in very low PA counseling rates.³⁰ Therefore, we have provided a simple stepwise process for clinicians prescribing PA to older adults with compromised function (Figure S2). For low-active older adults who present with difficulty ambulating, we suggest focusing first on developing strength and balance. The next *biggest return* recommendation would focus on the accumulation of more daily MVPA that can be done in brief bouts (eg, climbing a flight of stairs) to minimize task aversion. At this point, a shift in *moving more* via LPA is warranted. Finally, once individuals are doing a modest amount of PA across the full intensity continuum, one may focus on *moving more often*, allowing the individual to accumulate movement while guarding against the ill effects of sustained sitting.

Strengths and Limitations

This study leveraged a relatively large sample of older adults who had varied levels of compromised function recruited across a wide geographic area. We used a statistical approach that allowed for the modeling of the relationship between PA and MMD over time and included a lengthy follow-up period. A key limitation to this study was that the PA intervention was not designed to affect lifestyle LPA. As such, although we controlled for several baseline comorbidities, it may be that our analyses highlighted healthy individuals who were capable of engaging in more lifestyle movement. Future interventions attempting to directly manipulate both the amount and patterning of PA are warranted. Additionally, our model used the most recent accelerometer value to predict subsequent MMD risk. Due to having only three activity assessments, some predictions may be based on an individual's baseline PA levels alone. As a sensitivity analyses, we reran our models with only individuals with baseline and at least one follow-up assessment that reduced our sample size by 17 and had no meaningful effect on MMD risk. Fortunately, accelerometers are becoming easier to wear, and future research should consider replicating our analyses using more robust PA data sets.

In conclusion, PA is a vital health-promoting behavior for maintaining physical functioning and an independent lifestyle. For older adults at high risk for MMD, our findings underscore the importance of a tailored approach to PA promotion. At present, the necessary behavior change techniques that can effectively bring about a distributed pattern of movement are unknown, and we believe this represents an important path forward for research on promoting quality of life as individuals age.

ACKNOWLEDGMENTS

Financial Disclosure: This work was supported by the National Institutes of Health and National Institute on Aging (U01 AG22376), the National Heart, Lung and Blood Institute (3U01AG022376-05A2S), and the Wake Forest University Claude D. Pepper Older Americans Independence Center (P30-AG21332).

Conflict of Interest: The authors have declared no conflicts of interest for this article.

Author Contributions: Jason Fanning conceptualized the research question and led manuscript preparation. W. Jack Rejeski assisted in conceptualization and

manuscript preparation. Shyh-Huei Chen led data processing and management for the analyses and edited the manuscript. Michael E. Miller led data analysis and assisted in conceptualization and manuscript preparation. Jack Guralnik and Marco were involved in the conceptualization and conduct of the LIFE study and participated in manuscript preparation.

Sponsor's Role: The funding agencies did not participate in the design or conduct of the study; nor did they participate in the preparation of the manuscript at any stage.

REFERENCES

- Guralnik JM, Lacroix AZ, Abbott RD, et al. Maintaining mobility in late life. I. Demographic characteristics and chronic conditions. *Am J Epidemiol*. 1993;137(8):845-857.
- Loneragan ET, Krevans JR. A national agenda for research on aging. *N Engl J Med*. 1991;324(25):1825-1828.
- Buchner DM, Beresford SAA, Larson EB, LaCroix AZ, Wagner EH. Effects of physical activity on health status in older adults II: intervention studies. *Annu Rev Public Health*. 1992;13(1):469-488.
- Stuck AE, Walthert JM, Nikolaus T, Büla CJ, Hohmann C, Beck JC. Risk factors for functional status decline in community-living elderly people: a systematic literature review. *Soc Sci Med*. 1999;48(4):445-469.
- Warburton DER, Bredin SSD. Reflections on physical activity and health: what should we recommend? *Can J Cardiol*. 2016;32(12):1-10.
- Rejeski WJ, Axtell R, Fielding R, et al. Promoting physical activity for elders with compromised function: the Lifestyle Interventions and Independence for Elders (LIFE) study physical activity intervention. *Clin Interv Aging*. 2013;8:1119-1131.
- Rejeski WJ, Marsh AP, Brubaker PH, et al. Analysis and interpretation of accelerometry data in older adults: the LIFE study. *J Gerontol A Biol Sci Med Sci*. 2016;71(4):521-528.
- Pahor M, Guralnik JM, Ambrosius WT, et al. Effect of structured physical activity on prevention of major mobility disability in older adults: the LIFE study randomized clinical trial. *JAMA*. 2014;311(23):2387-2396.
- Rosenberg DE, Bombardier CH, Hoffman JM, Belza B. Physical activity among persons aging with mobility disabilities: shaping a research agenda. *J Aging Res*. 2011;2011:708510.
- McPhee JS, French DP, Jackson D, Nazroo J, Pendleton N, Degens H. Physical activity in older age: perspectives for healthy ageing and frailty. *Bio-gerontology*. 2016;17(3):567-580.
- Messier SP, Loeser RF, Mitchell MN, et al. Exercise and weight loss in obese older adults with knee osteoarthritis: a preliminary study. *J Am Geriatr Soc*. 2000;48(9):1062-1072.
- McDermott MM, Liu K, Guralnik JM, et al. Home-based walking exercise intervention in peripheral artery disease: a randomized clinical trial. *JAMA*. 2013;310(1):57-65.
- Rejeski WJ, Brubaker PH, Goff DC, et al. Translating weight loss and physical activity programs into the community to preserve mobility in older, obese adults in poor cardiovascular health. *Arch Intern Med*. 2011;171(10):880-886.
- Russell E. Exercise is medicine. *CMAJ*. 2013;185(11):E526.
- World Health Organization. New Physical Activity Recommendations for Reducing Disease and Prevent Deaths. http://www.who.int/chp/media/news/releases/2011_2_physicalactivity/en/. Accessed May 3, 2015.
- Fielding RA, Guralnik JM, King AC, et al. Dose of physical activity, physical functioning and disability risk in mobility-limited older adults: results from the LIFE study randomized trial. *PLOS One*. 2017;12(8):e0182155.
- Davis MG, Fox KR, Stathi A, Trayers T, Thompson JL, Cooper AR. Objectively measured sedentary time and its association with physical function in older adults. *J Aging Phys Act*. 2014;22:474-481.
- Fielding RA, Rejeski WJ, Blair S, et al. The lifestyle interventions and independence for elders study: design and methods. *J Gerontol A Biol Sci Med Sci*. 2011;66A(11):1226-1237.
- Borg GA. Perceived exertion: a note on "history" and methods. *Med Sci Sports*. 1973;5(2):90-93.
- Choi N. Relationship between health service use and health information technology use among older adults: analysis of the US National Health Interview Survey. *J Med Internet Res*. 2011;13(2):e33.
- Copeland JL, Eslinger DW. Accelerometer assessment of physical activity in active, healthy older adults. *J Aging Phys Act*. 2009;17(1):17-30. <http://europemc.org/abstract/med/19299836>. Accessed February 18, 2015.

22. Barnett A, Van Den Hoek D, Barnett D, Cerin E. Measuring moderate-intensity walking in older adults using the ActiGraph accelerometer. *BMC Geriatr.* 2016;16(1):211.
23. Rejeski WJ, Walkup MP, Fielding RA, LIFE Study Investigators. Evaluating accelerometry thresholds for detecting changes in levels of moderate physical activity and resulting major mobility disability. *J Gerontol A Biol Sci Med Sci.* 2017;73(5):660-667.
24. Fanning J, Rejeski WJ, Chen S-H, et al. A case for promoting movement medicine: preventing disability in the LIFE randomized controlled trial. *J Gerontol A Biol Sci Med Sci.* 2019;74(11):1821-1827.
25. Piercy KL, Troiano RP, Ballard RM, et al. The Physical Activity Guidelines for Americans. *JAMA.* 2018;320(19):2020-2028.
26. Hall KS, Howe CA, Rana SR, Martin CL, Morey MC. METs and accelerometry of walking in older adults: standard versus measured energy cost. *Med Sci Sports Exerc.* 2013;45(3):574-582.
27. National Health and Aging Trends Study. Public Use Data (Round 7). National Institute on Aging; Johns Hopkins Bloomberg School of Public Health; 2010 Accessed September 24, 2019.
28. Wanigatunga AA, Ferrucci L, Schrack JA. Physical activity fragmentation as a potential phenotype of accelerated aging. *Oncotarget.* 2019;10(8):807-809.
29. Marsh AP, Rejeski WJ, Espeland MA, et al. Muscle strength and BMI as predictors of major mobility disability in the Lifestyle Interventions and Independence for Elders pilot (LIFE-P). *J Gerontol A Biol Sci Med Sci.* 2011;66(12):1376-1383.
30. VanWormer JJ, Pronk NP, Kroeninger GJ. Clinical counseling for physical activity: translation of a systematic review into care recommendations. *Diabetes Spectr.* 2009;22(1):48-55.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article.

Figure S1: Utilizing 760 as the cut point for moderate to moderate to vigorous physical activity (MVPA), the difference in risk for major mobility disability (MMD) that a 1-unit higher level of variability has at different amounts of light physical activity (LPA) (vertical lines), and difference in risk associated with a 1-min/h higher level of amount at different levels of variability in LPA (horizontal lines) among those with a Short Physical Performance Battery (SPPB) score of 7 or lower (bottom panel) and SPPB score of 8 to 9, extracted from a model including all activity

predictors. Note that we have illustrated variability hazard ratios (HRs) at selected values of 6 vs 5, 9 vs 8, and 11 vs 10, and number of HRs at selected values of 7 vs 6 min/h, 13 vs 12 min/h, and 17 vs 16 min/h. Red dots represent activity measurement intervals followed by MMD events. HRs for LPA were obtained through back-transformation from model generated using log-transformed values of LPA.

Figure S2: Stepwise Activity Recommendation Guide. Recommendations build from left to right, such that fostering basic physical function is necessary for building a greater volume of higher intensity movement. Once higher intensity movement has been built into the day, and function is stable, a focus on lower intensity movement is warranted, followed by a focus on distribution of lower intensity movement. *It is vital to bear in mind that intensity is relevant, and as such, many basic activities of daily living (eg, walking for transport) should be seen as higher intensity for some individuals with very low levels of basic functioning.

Table S1: Development of Cox proportional hazards models using an accelerometer cut point of 1040 to define moderate to vigorous physical activity (MVPA). Each column represents a step in the development of the model for light physical activity (LPA) or MVPA, culminating in the final full model that retained all significant predictors identified in the model-building steps, plus covariates. BP = blood pressure, CVD = cardiovascular disease. HR = hazard ratio; SPPB = Short Physical Performance Battery.

Table S2: Development of Cox proportional hazards models using an accelerometer cut point of 760 to define moderate to vigorous physical activity (MVPA). Each column represents a step in the development of the model for light physical activity (LPA) or MVPA, culminating in the final full model that retained all significant predictors identified in the model-building steps, plus covariates. BP = blood pressure; CVD = cardiovascular disease; HR = hazard ratio; SPPB = Short Physical Performance Battery.